

Figure 3D-14.
 Simulated Brown Trout Fry Habitat in Segments 1-3 of Middle Owens River
 for the Alternatives, 1940-1989 Hydrologic Period




Table 3D-23.
 Average Brown Trout Fry Habitat in Segments 1-3 of Middle Owens River
 by Alternative, 1940-1989 Hydrologic Period

Alternative or Condition	Segment 1	Segment 2	Segment 3	Total
Average WUA (thousands of sq ft)				
Point-of-reference scenario	77,123	316,543	269,482	663,148
No restriction	77,989	313,501	271,342	662,833
6,372	75,935	322,466	269,895	668,296
6,377	75,786	326,723	270,254	672,763
6,383.5	74,871	331,400	268,172	674,444
6,390	74,412	333,414	268,072	675,898
6,410	74,394	336,627	266,129	677,150
No diversion	74,687	339,001	264,036	677,725
Percent Difference from Point-of-Reference Scenario				
No restriction	1.1	-1.0	0.7	-0.0
6,372	-1.5	1.9	0.2	0.8
6,377	-1.7	3.2	0.3	1.4
6,383.5	-2.9	4.7	-0.5	1.7
6,390	-3.5	5.3	-0.5	1.9
6,410	-3.5	6.3	-1.2	2.1
No diversion	-3.2	7.1	-2.0	2.2